



Inverness Dental 314 S Line Ave Inverness, FL 34452 P: 352-419-4056 F: 888-411-2260	Inverness Medical 300 S Line Ave, Inverness, FL 34452 P: 352-419-5760 F: 877-209-5265	Lecanto Medical 595 N Lecanto Hwy Lecanto, FL 34461 P: 352-527-2244 F: 877-275-8781	Ocala 7205 SE Maricamp Rd Ocala, FL 34472 P: 352-680-7000 F: 877-849-9264	Sumterville 1389 S US Hwy 301 Sumterville, FL 33585 P: 352-793-5900 F: 855-832-3504	Homosassa 7945 S Suncoast Blvd Suites A & B Homosassa, FL 34446 P: 352-423-4923 F: 352-423-1468
--	--	--	--	--	---

Hi Parent/Guardian,

Langley Health Services is pleased to be coordinating with the Sumter County School Board to offer vaccines! We will be at your child's school on the following dates:

School	Time of day	Date
South Sumter Middle	9:30am – 10:30am	4/11/2024
Sumter PREP Academy	11:00am – 11:30am	4/11/2024
Wildwood Intermediate School	9:30am – 10:30am	4/16/2024

Vaccines to be offered include Tdap (tetanus, diphtheria, acellular pertussis), meningococcal ACWY, Hepatitis A, and HPV. The Tdap vaccine is required for all students to enter 7th grade. If you would like for your child to receive any of the vaccines offered, please complete the attached consent form. On the form, in the block under the vaccine name, you will need to initial for each vaccine you consent for your child to receive. You will return this form to your child's school. Please return the form no later than Friday, April 5, 2024 for South Sumter Middle School and Sumter PREP Academy and Friday, April 12, 2024 for Wildwood Intermediate School.

For your future healthcare needs, keep in mind we offer the following services: medical (including Family Practice, Pediatrics, and walk-in), dental, behavioral health, radiology (x-rays, MRIs, CT scans, ultrasounds, etc.), podiatry (foot and ankle care), and pharmacy services. We have locations in Sumter, Citrus, and Marion counties. If you have any questions, or need any assistance, please let us know! We will be happy to help you.

Thank you,
Your Langley Health Services team

Homeroom teacher: _____



School Vaccine Consent Form

Please write legibly and complete all blank fields.

Student's Name (last, first, middle initial): _____			
Student's Date of Birth: _____		Student's Social Security Number: _____	
Parent/Guardian Phone Number: () _____		Parent/guardian Email: _____	
Student's Address: _____			
Student's Sex (gender assigned at birth): <input type="checkbox"/> Female <input type="checkbox"/> Male		Student's Ethnicity: <input type="checkbox"/> Hispanic/Latino <input type="checkbox"/> Not Hispanic/Latino <input type="checkbox"/> Unknown	
Student's Race: <div style="display: flex; justify-content: space-between;"> <div> <input type="checkbox"/> American Indian/Alaskan Native <input type="checkbox"/> Black/African American <input type="checkbox"/> Other Pacific Islander <input type="checkbox"/> Unknown </div> <div> <input type="checkbox"/> Asian <input type="checkbox"/> Native Hawaiian <input type="checkbox"/> White </div> </div>			
Student's Insurance screening: Please complete this section entirely. <input type="checkbox"/> Medicaid <input type="checkbox"/> No health insurance <input type="checkbox"/> Insured <i>If student is insured:</i> Primary Insurance Carrier ID #: _____ Grp #: _____ Insurance Company: _____ Insurance Company Phone #: _____ Insured's Name: _____ Insured's Relationship to Patient: _____ Insured's Date of Birth: _____ Secondary Insurance Carrier ID #: _____ Grp #: _____ Insurance Company: _____ Insurance Company Phone #: _____ Insured's Name: _____ Insured's Relationship to Patient: _____ Insured's Date of Birth: _____			
STUDENT'S SCHOOL INFORMATION-			
Which school does the student attend? _____			
What grade is the student in? _____			
Who is the student's homeroom teacher? _____			
VACCINE HEALTH SCREENING- Please answer all questions about the student who will be receiving the vaccine(s). You will answer by placing a check mark in the correct "yes" or "no" column. Answers will determine whether the student can be vaccinated at this time. YES NO			
Is the student to be vaccinated sick today? <i>Temperature will be taken prior to vaccination.</i>			
Does the student have any allergies to medication, foods, latex, or any vaccines? If yes, please explain the allergy: _____			
Has the person to be vaccinated ever had a serious reaction to Tdap vaccine in the past?			
Does the child have a long-term health problem with lung, heart, kidney or metabolic disease (e.g., diabetes), asthma, a blood disorder, no spleen, complement component deficiency, a cochlear implant, or a spinal fluid leak?			
Has the child received vaccinations in the past 4 weeks?			

CONSENT TO VACCINATE

I (parent or guardian of student named above) have been given a copy and I have read, or had explained to me, the information in the Vaccine Information Statement(s) for the vaccine(s) my child will be receiving. I understand that for coordination purposes only, this information may be shared with the staff at my child's school and/or the Sumter County School Board. I have had a chance to ask questions and fully understand the benefits and risks of each of the indicated vaccines and ask the following vaccine be given to my child on the scheduled school clinic date (**initial all that apply**):

Tetanus, diphtheria, acellular pertussis (Tdap) <i>Tetanus-diphtheria-pertussis (Tdap) immunization is required for all students entering 7th grade</i>	Meningococcal ACWY (MCV4)	Hepatitis A	HPV

Printed Name of Parent or Guardian of Patient: _____	Signature of Parent or Guardian of Patient: _____	Today's Date: _____
--	---	---------------------

Tdap (Tetanus, Diphtheria, Pertussis) Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Tdap vaccine can prevent **tetanus, diphtheria, and pertussis**.

Diphtheria and pertussis spread from person to person. Tetanus enters the body through cuts or wounds.

- **TETANUS (T)** causes painful stiffening of the muscles. Tetanus can lead to serious health problems, including being unable to open the mouth, having trouble swallowing and breathing, or death.
- **DIPHTHERIA (D)** can lead to difficulty breathing, heart failure, paralysis, or death.
- **PERTUSSIS (aP)**, also known as “whooping cough,” can cause uncontrollable, violent coughing that makes it hard to breathe, eat, or drink. Pertussis can be extremely serious especially in babies and young children, causing pneumonia, convulsions, brain damage, or death. In teens and adults, it can cause weight loss, loss of bladder control, passing out, and rib fractures from severe coughing.

2. Tdap vaccine

Tdap is only for children 7 years and older, adolescents, and adults.

Adolescents should receive a single dose of Tdap, preferably at age 11 or 12 years.

Pregnant people should get a dose of Tdap during every pregnancy, preferably during the early part of the third trimester, to help protect the newborn from pertussis. Infants are most at risk for severe, life-threatening complications from pertussis.

Adults who have never received Tdap should get a dose of Tdap.

Also, **adults should receive a booster dose of either Tdap or Td** (a different vaccine that protects against tetanus and diphtheria but not pertussis) **every 10 years**, or after 5 years in the case of a severe or dirty wound or burn.

Tdap may be given at the same time as other vaccines.

3. Talk with your health care provider


Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any vaccine that protects against tetanus, diphtheria, or pertussis**, or has any **severe, life-threatening allergies**
- Has had a **coma, decreased level of consciousness, or prolonged seizures within 7 days after a previous dose of any pertussis vaccine (DTP, DTaP, or Tdap)**
- Has **seizures or another nervous system problem**
- Has ever had **Guillain-Barré Syndrome** (also called “GBS”)
- Has had **severe pain or swelling after a previous dose of any vaccine that protects against tetanus or diphtheria**

In some cases, your health care provider may decide to postpone Tdap vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting Tdap vaccine.

Your health care provider can give you more information.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

4. Risks of a vaccine reaction

- Pain, redness, or swelling where the shot was given, mild fever, headache, feeling tired, and nausea, vomiting, diarrhea, or stomachache sometimes happen after Tdap vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines.



Meningococcal ACWY Vaccine:

What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Meningococcal ACWY vaccine can help protect against **meningococcal disease** caused by serogroups A, C, W, and Y. A different meningococcal vaccine is available that can help protect against serogroup B.

Meningococcal disease can cause meningitis (infection of the lining of the brain and spinal cord) and infections of the blood. Even when it is treated, meningococcal disease kills 10 to 15 infected people out of 100. And of those who survive, about 10 to 20 out of every 100 will suffer disabilities such as hearing loss, brain damage, kidney damage, loss of limbs, nervous system problems, or severe scars from skin grafts.

Meningococcal disease is rare and has declined in the United States since the 1990s. However, it is a severe disease with a significant risk of death or lasting disabilities in people who get it.

Anyone can get meningococcal disease. Certain people are at increased risk, including:

- Infants younger than one year old
- Adolescents and young adults 16 through 23 years old
- People with certain medical conditions that affect the immune system
- Microbiologists who routinely work with isolates of *N. meningitidis*, the bacteria that cause meningococcal disease
- People at risk because of an outbreak in their community


2. Meningococcal ACWY vaccine

Adolescents need 2 doses of a meningococcal ACWY vaccine:

- First dose: 11 or 12 years of age
- Second (booster) dose: 16 years of age

In addition to routine vaccination for adolescents, meningococcal ACWY vaccine is also recommended for **certain groups of people**:

- People at risk because of a serogroup A, C, W, or Y meningococcal disease outbreak
- People with HIV
- Anyone whose spleen is damaged or has been removed, including people with sickle cell disease
- Anyone with a rare immune system condition called “complement component deficiency”
- Anyone taking a type of drug called a “complement inhibitor,” such as eculizumab (also called “Soliris”®) or ravulizumab (also called “Ultomiris”®)
- Microbiologists who routinely work with isolates of *N. meningitidis*
- Anyone traveling to or living in a part of the world where meningococcal disease is common, such as parts of Africa
- College freshmen living in residence halls who have not been completely vaccinated with meningococcal ACWY vaccine
- U.S. military recruits



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of meningococcal ACWY vaccine**, or has any **severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone meningococcal ACWY vaccination until a future visit.

There is limited information on the risks of this vaccine for pregnant or breastfeeding people, but no safety concerns have been identified. A pregnant or breastfeeding person should be vaccinated if indicated.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting meningococcal ACWY vaccine.

Your health care provider can give you more information.

4. Risks of a vaccine reaction

- Redness or soreness where the shot is given can happen after meningococcal ACWY vaccination.
- A small percentage of people who receive meningococcal ACWY vaccine experience muscle pain, headache, or tiredness.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines.



HPV (Human Papillomavirus) Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

HPV (human papillomavirus) vaccine can prevent infection with some types of human papillomavirus.

HPV infections can cause certain types of cancers, including:

- cervical, vaginal, and vulvar cancers in women
- penile cancer in men
- anal cancers in both men and women
- cancers of tonsils, base of tongue, and back of throat (oropharyngeal cancer) in both men and women

HPV infections can also cause anogenital warts.

HPV vaccine can prevent over 90% of cancers caused by HPV.

HPV is spread through intimate skin-to-skin or sexual contact. HPV infections are so common that nearly all people will get at least one type of HPV at some time in their lives. Most HPV infections go away on their own within 2 years. But sometimes HPV infections will last longer and can cause cancers later in life.

2. HPV vaccine

HPV vaccine is routinely recommended for adolescents at 11 or 12 years of age to ensure they are protected before they are exposed to the virus. HPV vaccine may be given beginning at age 9 years and vaccination is recommended for everyone through 26 years of age.

HPV vaccine may be given to adults 27 through 45 years of age, based on discussions between the patient and health care provider.

Most children who get the first dose before 15 years of age need 2 doses of HPV vaccine. People who get the first dose at or after 15 years of age and younger people with certain immunocompromising conditions need 3 doses. Your health care provider can give you more information.

HPV vaccine may be given at the same time as other vaccines.

3. Talk with your health care provider


Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of HPV vaccine**, or has any **severe, life-threatening allergies**
- Is **pregnant**—HPV vaccine is not recommended until after pregnancy

In some cases, your health care provider may decide to postpone HPV vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting HPV vaccine.

Your health care provider can give you more information.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

4. Risks of a vaccine reaction

- Soreness, redness, or swelling where the shot is given can happen after HPV vaccination.
- Fever or headache can happen after HPV vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines.



Hepatitis A Vaccine:

What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Hepatitis A vaccine can prevent **hepatitis A**.

Hepatitis A is a serious liver disease. It is usually spread through close, personal contact with an infected person or when a person unknowingly ingests the virus from objects, food, or drinks that are contaminated by small amounts of stool (poop) from an infected person.

Most adults with hepatitis A have symptoms, including fatigue, low appetite, stomach pain, nausea, and jaundice (yellow skin or eyes, dark urine, light-colored bowel movements). Most children less than 6 years of age do not have symptoms.

A person infected with hepatitis A can transmit the disease to other people even if he or she does not have any symptoms of the disease.

Most people who get hepatitis A feel sick for several weeks, but they usually recover completely and do not have lasting liver damage. In rare cases, hepatitis A can cause liver failure and death; this is more common in people older than 50 years and in people with other liver diseases.

Hepatitis A vaccine has made this disease much less common in the United States. However, outbreaks of hepatitis A among unvaccinated people still happen.

2. Hepatitis A vaccine

Children need 2 doses of hepatitis A vaccine:

- First dose: 12 through 23 months of age
- Second dose: at least 6 months after the first dose

Infants 6 through 11 months old traveling outside the United States when protection against hepatitis A is recommended should receive 1 dose of hepatitis A vaccine. These children should still get 2 additional doses at the recommended ages for long-lasting protection.

Older children and adolescents 2 through 18 years of age who were not vaccinated previously should be vaccinated.


Adults who were not vaccinated previously and want to be protected against hepatitis A can also get the vaccine.

Hepatitis A vaccine is also recommended for the following people:

- International travelers
- Men who have sexual contact with other men
- People who use injection or non-injection drugs
- People who have occupational risk for infection
- People who anticipate close contact with an international adoptee
- People experiencing homelessness
- People with HIV
- People with chronic liver disease

In addition, a person who has not previously received hepatitis A vaccine and who has direct contact with someone with hepatitis A should get hepatitis A vaccine as soon as possible and within 2 weeks after exposure.

Hepatitis A vaccine may be given at the same time as other vaccines.



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of hepatitis A vaccine**, or has any **severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone hepatitis A vaccination until a future visit.

Pregnant or breastfeeding people should be vaccinated if they are at risk for getting hepatitis A. Pregnancy or breastfeeding are not reasons to avoid hepatitis A vaccination.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting hepatitis A vaccine.

Your health care provider can give you more information.

4. Risks of a vaccine reaction

- Soreness or redness where the shot is given, fever, headache, tiredness, or loss of appetite can happen after hepatitis A vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines.

